

Infrared regular representation of the three dimensional massless Nelson model

József Lőrinczi¹, Robert A. Minlos² and Herbert Spohn¹

¹*Zentrum Mathematik, Technische Universität München
80290 München, Germany
lorinczi@mathematik.tu-muenchen.de
spohn@mathematik.tu-muenchen.de*

²*Dobrushin Mathematics Laboratory
Institute for Information Transmission Problems
Bolshoy Karetny per. 19, 101447 Moscow, Russia
minl@iitp.ru*

Abstract

We prove that in the Euclidean representation of the three dimensional massless Nelson model the $t = 0$ projection of the interacting measure is absolutely continuous with respect to a Gaussian measure with suitably adjusted mean. We also determine the Hamiltonian in the Fock space over this Gaussian measure space.

KEYWORDS: Nelson's scalar field model, infrared regular representation, ground state, Gibbs measure on path space, cluster expansion

1 Introduction

The Nelson model [6] of a spinless electron coupled to a scalar massless Bose field is infrared divergent in 3 space dimensions. In its algebraic version the ground state representation of the Nelson Hamiltonian is not unitarily equivalent to the Fock representation [3, 2]. From the point of view of functional integrals, which is the one taken here, the $t = 0$ projection of the interacting measure is singular with respect to the $t = 0$ projection of the free measure [5]. Such a result leaves open the possibility of an explicit representation of the $t = 0$ functional measure. In this letter we will settle the issue. In the last section we will explain the connection to the non-Fock representation.

The infrared divergence in Nelson's model can be seen already on the level of the classical action. The Euclidean action for the electron is

$$S_p(\{q_t\}) = \int_{-\infty}^{\infty} \left(\frac{1}{2} \dot{q}_t^2 + V(q_t) \right) dt. \quad (1.1)$$

Here $t \mapsto q_t \in \mathbb{R}^3$ is a path of the electron. The electron's mass is set equal to one only for convenience. $V : \mathbb{R}^3 \rightarrow \mathbb{R}$ is a confining potential, which for technical reasons we require to be bounded from below, continuous and have the asymptotics $V(x) = C|x|^{2s} + o(|x|^{2s})$ for large $|x|$ with some constant $C > 0$ and exponent $s > 1$. The massless field $\xi_t : \mathbb{R}^3 \rightarrow \mathbb{R}$ has the action

$$S_f(\{\xi_t\}) = \int_{-\infty}^{\infty} \int_{\mathbb{R}^3} \frac{1}{2} (\partial_t \xi_t(x)^2 + \nabla \xi_t(x)^2) dx dt. \quad (1.2)$$

Finally, the electron is coupled locally and translation invariant as given by

$$S_{\text{int}}(\{q_t, \xi_t\}) = \int_{-\infty}^{\infty} \int_{\mathbb{R}^3} \varrho(x - q_t) \xi_t(x) dx dt. \quad (1.3)$$

Here ϱ is a form factor, assumed to be a sufficiently smooth function and of rapid decrease, normalized such that $\int_{\mathbb{R}^3} \varrho(x) dx = e$, where e is the charge of the particle. The case of interest is $e \neq 0$, since $e = 0$ implies $\hat{\varrho}(0) = 0$ ($\hat{\varrho}$ the Fourier transform of ϱ) and therefore regularizes the interaction in the infrared region. At a later stage we will need that $|e| \leq e^*$ with some sufficiently small e^* .

The infrared divergence can be deduced from the solution of the classical field equation which minimizes energy. An easy computation leads to the minimizer $q_t = q_{\min}$, where $q_{\min}$ corresponds to an absolute minimum of the potential, i.e. $V(x) \geq V(q_{\min})$, for all $x \in \mathbb{R}^3$, and $\xi_t = \xi_{\min}$ given by

$$\xi_{\min}(x) = \Delta^{-1} \varrho(x - q_{\min}). \quad (1.4)$$

In three dimensions $\xi_{\min}(x) \cong -e/(4\pi|x|)$ for large $|x|$. On the other hand, for the free measure $\varrho = 0$ and $\xi_{\min} = 0$. If the total action $S = S_p + S_f + S_{\text{int}}$ is used for constructing the Euclidean path measure (formally like $\exp(-S)$), then basically one has to compare two Gaussian fields of the same covariance, one with mean $\Delta^{-1} \varrho(x)$ and the other with zero mean. Because of slow decay of the Coulomb potential in 3 dimensions, these two measures cannot have a density with respect to each other.

Our arguments also offer a hint of how to improve on this situation. We reshuffle the field action by adding a term dependent on a function u to be chosen below:

$$S_f^u(\{\xi_t\}) = \int_{-\infty}^{\infty} \int_{\mathbb{R}^3} \left(\frac{1}{2} (\partial_t \xi_t(x)^2 + (\nabla \xi_t(x))^2) + \xi_t(x)u(x) \right) dx dt. \quad (1.5)$$

For convenience, we represent this auxiliary function as the convolution

$$u(x) = (\varrho * h)(x) \equiv \int_{\mathbb{R}^3} e^{ik \cdot x} \hat{\varrho}(k) \hat{h}(k) dk \quad (1.6)$$

where $\hat{h}$ denotes the Fourier transform of h , of which we require that

1. $\hat{h}$ is a real-valued, even, bounded, and sufficiently smooth function;
2. $\hat{h}(0) = 1$.

The first condition is more for simplicity than necessity, the second is essential. The density $\exp(-S_f^u(\{\xi_t\}))$ provides the free Gaussian measure. Keeping the electron's action and the total action unchanged, we obtain a reshuffled interaction of the form

$$S_{\text{int}}^u(\{q_t, \xi_t\}) = \int_{-\infty}^{\infty} \int_{\mathbb{R}^3} (\varrho(x - q_t) - (\varrho * h)(x)) \xi_t(x) dx dt. \quad (1.7)$$

The function u is chosen in such a way that the free Gaussian's mean asymptotically agrees with the solution of minimal energy of the classical field equation. This then makes up for the possibility of the $t = 0$ interacting measure to be absolutely continuous with respect to the $t = 0$ projection of the above Gaussian measure. In the remainder of the letter this argument will be made sound.

2 Infrared regular representation

First we translate the actions to well defined measures on path space. S_p is associated with the stationary $P(\phi)_1$ -process q_t on $C(\mathbb{R}, \mathbb{R}^3)$ defined by the stochastic differential equation

$$dq_t = (\nabla \log \psi_0) dt + dB_t \quad (2.1)$$

(dB_t denotes Brownian motion), with invariant measure $d\mathbf{N}^0 = \psi_0^2(x) dx$. Here ψ_0 is the ground state of the Schrödinger operator $(-1/2)\Delta + V(q)$ generating the $P(\phi)_1$ -process. The path measure of this process will be denoted by $\mathcal{N}^0$. The paths $Q = \{q_t : t \in \mathbb{R}\}$ are $\mathcal{N}^0$ -almost surely continuous and q_t is a time-reversible Markov process. Moreover, for our class of potentials V the semigroup of this process is intrinsically ultracontractive [4].

The field $\{\xi_t(f) : f \in \mathcal{S}(\mathbb{R}^3), t \in \mathbb{R}\}$ with $f \in \mathcal{S}(\mathbb{R}^3)$, i.e. the Schwartz space over $\mathbb{R}^3$, is in its turn described by the Gaussian measure $\mathcal{G}$ on $C(\mathbb{R}, \mathcal{S}'(\mathbb{R}^3))$ resulting from the (modified) Euclidean action S_f^u . This Gaussian measure has mean

$$\mathbb{E}_{\mathcal{G}}[\xi_t(f)] = - \int_{\mathbb{R}^3} \frac{\hat{\varrho}(k) \hat{f}(k) \hat{h}(k)}{|k|^2} dk \equiv \int_{\mathbb{R}^3} f(x) \gamma(x) dx \quad (2.2)$$

(i.e., represented in position space in terms of the function γ), and covariance

$$\text{cov}_{\mathcal{G}}(\xi_s(f_1), \xi_t(f_2)) = \int_{\mathbb{R}^3} \frac{\hat{f}_1(k) \hat{f}_2^*(k)}{2|k|} e^{-|k||t-s|} dk. \quad (2.3)$$

We denote the $t = 0$ stationary distribution of this Markov process by $\mathbf{G}$; it is itself a Gaussian measure with mean (2.2) and covariance (2.3) taken with $s = t$. For easing the argument below, we introduce the centred field

$$\eta_t(f) = \xi_t(f) - \mathbb{E}_{\mathcal{G}}[\xi_t(f)], \quad f \in \mathcal{S}(\mathbb{R}^3); \quad (2.4)$$

as an auxiliary variable; this obviously has then a Gaussian distribution with zero mean and the same covariance as (2.3) above. Note that u introduces a shift in the mean of the unmodified Gaussian measure $\mathcal{G}_0$ on the same space (which with $h \equiv 0$ would be 0). We denote by $\mathbf{G}_0$ the $t = 0$ distribution of $\mathcal{G}_0$. This shift induces the unitary map

$$\mathcal{U} : L^2(\mathcal{S}'(\mathbb{R}^3), d\mathbf{G}) \rightarrow L^2(\mathcal{S}'(\mathbb{R}^3), d\mathbf{G}_0), \quad (\mathcal{U}F)(f) = F(f - \gamma), \quad (2.5)$$

with $F : \mathcal{S}'(\mathbb{R}^3) \rightarrow \mathbb{R}$, $f \in \mathcal{S}'(\mathbb{R}^3)$ and γ given by (2.2), which we introduce here for later use.

The non-interacting joint particle-field process is set up on $C(\mathbb{R}, \mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3))$ and described by the path measure $\mathcal{P}^0 = \mathcal{N}^0 \times \mathcal{G}$ with $t = 0$ distribution $\mathbf{P}^0 = \mathbf{N}^0 \times \mathbf{G}$. The interacting system can be described by taking $\mathcal{P}^0$ as reference process and modifying it with the density given by

$$\frac{d\mathcal{P}_T}{d\mathcal{P}^0}(q, \xi) = \frac{1}{Z_T} \exp \left(- \int_{-T}^T S_{\text{int}}^u(\{q_t, \xi_t\}) dt \right) \quad (2.6)$$

Here

$$Z_T = \int \exp \left(- \int_{-T}^T S_{\text{int}}^u(\{q_t, \xi_t\}) dt \right) d\mathcal{P}^0 \quad (2.7)$$

is the normalizing partition function. To have a more explicit formula instead of (2.7) we first make the Gaussian part of the integration to obtain

$$\begin{aligned} Z_T &= \mathbb{E}_{\mathcal{N}_T^0} \left[\mathbb{E}_{\mathcal{G}} \left[\exp \left(- \int_{-T}^T \int_{\mathbb{R}^3} \hat{\xi}_t(k) \hat{\rho}(k) (e^{ik \cdot q_t} - \hat{h}(k)) dk dt \right) \right] \right] \\ &= \mathbb{E}_{\mathcal{N}_T^0} [Z_T^Q] \end{aligned}$$

where

$$\begin{aligned} Z_T^Q &= \int \exp \left(- \int_{-T}^T \int_{\mathbb{R}^3} \hat{\eta}_t(k) \hat{\rho}(k) (e^{ik \cdot q_t} - \hat{h}(k)) dk dt \right) \times \\ &\quad \times \exp \left(\int_{-T}^T \int_{\mathbb{R}^3} \hat{\gamma}(k) \hat{\rho}(k) (e^{ik \cdot q_t} - \hat{h}(k)) dk dt \right) d\mathcal{G}_0 \\ &= \exp \left(- \frac{1}{2} \int_{\mathbb{R}^3} \frac{|\hat{\rho}(k)|^2 \hat{h}(k)}{|k|} dk \int_{-\infty}^{\infty} ds \int_{-T}^T dt e^{-|k||t-s|} (e^{ik \cdot q_t} - \hat{h}(k)) \right) \times \\ &\quad \times \exp \left(\frac{1}{4} \int_{-T}^T \int_{-T}^T \frac{|\hat{\rho}(k)|^2}{|k|} dk (e^{ik \cdot q_s} - \hat{h}(k)) (e^{-ik \cdot q_t} - \hat{h}(k)) e^{-|k||t-s|} dt ds \right). \end{aligned} \quad (2.8)$$

Notice that in the above expression we passed from integration with respect to $\mathcal{G}$ to integration with respect to $\mathcal{G}_0$. From this we get that

$$Z_T^Q = C_T Z_T^Q \quad (2.9)$$

with $C_T > 0$ independent of Q and

$$\begin{aligned} Z_T^Q = & \exp \left(- \int_{-T}^T \int_{-T}^T W(q_s - q_t, s - t) dt ds \right) \times \\ & \exp \left(- \int_{-T}^T dt \int_{|s|>T} ds \int_{\mathbb{R}^3} W(q_t - q, t - s) h(q) dq \right) \times \\ & \times \exp \left(- \int_{-T}^T ds \int_{|t|>T} dt \int_{\mathbb{R}^3} W(q_s - q, t - s) h(q) dq \right). \end{aligned} \quad (2.10)$$

Here

$$W(q, t) = -\frac{1}{4} \int_{\mathbb{R}^3} \frac{|\hat{\varrho}(k)|^2}{|k|} e^{-|k||t|} \cos(k \cdot q) dk \quad (2.11)$$

This implies that the marginal distribution $\mathcal{N}_T$ of $\mathcal{P}_T$ for the particle paths satisfies

$$\frac{d\mathcal{N}_T}{d\mathcal{N}^0} = \frac{Z_T^Q}{Z_T}. \quad (2.12)$$

with $Z_T = \int Z_T^Q d\mathcal{N}^0$. The $t = 0$ single time distribution of $\mathcal{N}_T$ will be denoted by $\mathbf{N}_T$.

Lemma 2.1 *There is some $e^* > 0$ such that for all $|e| \leq e^*$ the weak local limit $\lim_{T \rightarrow \infty} \mathcal{N}_T = \mathcal{N}$ exists and is a Gibbs probability measure on $C(\mathbb{R}, \mathbb{R}^3)$. Moreover, $\mathcal{N}$ does not depend on h , i.e. it coincides with the limiting measure constructed for $h \equiv 0$.*

Proof: By using cluster expansion, the Gibbs measure for $h \equiv 0$ can be shown to exist for W at sufficiently weak couplings $|e| \leq e^*$. Here we have an interaction which differs from the cases studied in [4] by the last two factors in (2.10). However, they appear as extra energies coming from interactions between pieces of paths lying inside $[-T, T]$ with constant boundary conditions in $\mathbb{R} \setminus [-T, T]$ weighted by h , to which our previous arguments extend directly. $\square$

For any fixed path $Q = \{q_t : t \in \mathbb{R}\} \in C(\mathbb{R}, \mathbb{R}^3)$ consider now the conditional distribution $\mathcal{P}_T^Q = \mathcal{P}_T(\cdot | \{q_t\} = Q)$ as a probability measure on $C(\mathbb{R}, \mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3))$. Since the interaction is linear in ξ_t , $\mathcal{P}_T^Q$ is itself a Gaussian measure with covariance (2.3) and mean

$$\mathbb{E}_{\mathcal{P}_T^Q}[\xi_t(f)] = \int_{\mathbb{R}^3} \hat{f}(k) \hat{g}_T^{Q,t}(k) dk \quad (2.13)$$

where

$$\begin{aligned} \hat{g}_T^{Q,t}(k) &= \hat{\gamma}(k) - \frac{\hat{\varrho}(k)}{2|k|} \int_{-T}^T e^{-|k||t-\tau|} \left(e^{ik \cdot q_\tau} - \hat{h}(k) \right) d\tau \\ &= -\frac{\hat{\varrho}(k)}{2|k|} \int_{-T}^T e^{-|k||t-\tau|} e^{ik \cdot q_\tau} d\tau - \frac{\hat{\varrho}(k) \hat{h}(k)}{2|k|} \int_{|\tau|>T} e^{-|k||t-\tau|} d\tau. \end{aligned} \quad (2.14)$$

Lemma 2.2 *There is some $e^* > 0$ such that for all $|e| \leq e^*$ the weak local limit $\lim_{T \rightarrow \infty} \mathcal{P}_T = \mathcal{P}$ exists and is a probability measure on $C(\mathbb{R}, \mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3))$. Moreover, $\mathcal{P}$ coincides with the path measure for $h \equiv 0$.*

Proof: By using (2.14) it is immediate that $\lim_{T \rightarrow \infty} \hat{g}_T^{Q,t} = \hat{g}^{Q,t}$ exists and coincides with the mean for $\mathcal{P}^Q$ with $h \equiv 0$. Thus we have also that $\mathcal{P}_T^Q \rightarrow \mathcal{P}^Q$ exists. Moreover, since $\mathcal{P}_T(E \times F) = \int_F \mathcal{P}_T^Q(E) d\mathcal{N}_T$, for all $T > 0$, where E and F are appropriate measurable sets, we also have the claim of the lemma. Hence the joint particle-field path measure does not depend on h , i.e. on how the reshuffling of the free field and interaction energies is performed. $\square$

$\mathcal{P}$ and $\mathcal{P}^0$ are Markov processes whose $t = 0$ distributions we denote below by $\mathbf{P}$ resp. $\mathbf{P}^0$.

In the modified function space representation Nelson's Hamiltonian is H on the Hilbert space $\mathcal{H}^0 = L^2(\mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3), d\mathbf{P}^0)$ acting like

$$(F, e^{-tH}G)_{\mathcal{H}^0} = \mathbb{E}_{\mathbf{P}^0} \left[F(q_0, \xi_0) G(q_t, \xi_t) \exp\left(-\int_0^t \mathcal{L}_{\text{int}}^u(\{q_s, \xi_s\}) ds\right) \right], \quad t > 0 \quad (2.15)$$

with suitable F and G , and where $\mathcal{L}_{\text{int}}^u(\{q_t, \xi_t\}) = \int_{\mathbb{R}^3} (\varrho(x - q_t) - (\varrho * h)(x)) \xi_t(x) dx$ is the Lagrangian corresponding to (1.7). The bottom of the spectrum of H will be denoted by E_0 . We define the Euclidean Hamiltonian H_{euc} acting on the Hilbert space $\mathcal{H} = L^2(\mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3), d\mathbf{P})$ as the self-adjoint operator generating the symmetric contracting semigroup T_t given by

$$(F, T_t G)_{\mathcal{H}} = \mathbb{E}_{\mathcal{P}}[F(q_0, \xi_0) G(\xi_t, q_t)], \quad t > 0. \quad (2.16)$$

Theorem 2.3 *Suppose $|e| \leq e^*$ with some $e^* > 0$. Then H has a unique strictly positive ground state Ψ_0 corresponding to the lowest eigenvalue E_0 , and $H - E_0$ is unitarily equivalent with H_{euc} .*

Proof: From [5] we know that the existence of a strictly positive ground state of H implies that $H - E_0$ and H_{euc} are unitarily equivalent. Moreover, by Theorem 4.1 of the same reference we also know that for proving existence of the ground state it suffices to show that

$$\liminf_{T \rightarrow \infty} \int \left(\frac{d\mathbf{P}_T}{d\mathbf{P}^0} \right)^{1/2} d\mathbf{P}^0 > 0. \quad (2.17)$$

We have

$$\begin{aligned} \frac{d\mathbf{P}_T^Q}{d\mathbf{G}}(\xi) &= \exp \left(\int_{\mathbb{R}^3} (\hat{\xi}(k) - \hat{\gamma}(k)) \hat{\varrho}(k) dk \int_{-T}^T (e^{ik \cdot q_t} - \hat{h}(k)) e^{-|k||t|} dt \right) \times \\ &\exp \left(-\frac{1}{4} \int_{\mathbb{R}^3} dk \int_{-T}^T \int_{-T}^T \frac{|\hat{\varrho}(k)|^2}{|k|} (e^{ik \cdot q_t} - \hat{h}(k)) (e^{-ik \cdot q_s} - \hat{h}(k)) e^{-|k|(|t|+|s|)} dt ds \right). \end{aligned}$$

Moreover,

$$\frac{d\mathbf{P}_T}{d\mathbf{P}^0}(q, \xi) = \mathbb{E}_{\mathcal{N}_T} \left[\frac{d\mathbf{P}_T^Q}{d\mathbf{G}}(\xi) | q_0 = q \right] \frac{d\mathbf{N}_T}{d\mathbf{N}^0}(q). \quad (2.18)$$

By using that for some $c > 0$

$$\frac{1}{c} \leq \frac{d\mathbf{N}_T}{d\mathbf{N}^0} \leq c \quad (2.19)$$

obtained similarly as in [4], in combination with Jensen's inequality we obtain

$$\begin{aligned} & \int \left(\frac{d\mathbf{P}_T}{d\mathbf{P}^0}(q, \xi) \right)^{1/2} d\mathbf{P}^0(q) \\ &= \int \mathbb{E}_{\mathcal{N}_T} \left[\left(\frac{d\mathbf{P}^Q}{d\mathbf{G}} \right)^{1/2} \middle| q = q_0 \right] \left(\frac{d\mathbf{N}_T}{d\mathbf{N}^0}(q) \right)^{1/2} d\mathbf{P}^0 \\ &\geq \frac{1}{\sqrt{c}} \int \exp \left(\mathbb{E}_{\mathcal{N}_T} \left[\frac{1}{2} \int_{\mathbb{R}^3} (\hat{\xi}(k) - \hat{\gamma}(k)) \hat{\rho}(k) \int_{-T}^T e^{-|k||t|} (e^{ik \cdot q_t} - \hat{h}(k)) dt dk \middle| q = q_0 \right] \right) \\ &\quad \times \exp \left(-\frac{1}{8} \int_{\mathbb{R}^3} dk \int_{-T}^T \int_{-T}^T \frac{|\hat{\rho}(k)|^2}{|k|} m_T(k; s, t) e^{-|k|(|t|+|s|)} dt ds \right) d\mathbf{P}^0 \\ &\geq \frac{1}{\sqrt{c}} \exp \left(-\frac{c}{8} \int_{\mathbb{R}^3} dk \int_{-T}^T \int_{-T}^T \frac{|\hat{\rho}(k)|^2}{|k|} m_T(k; s, t) e^{-|k|(|t|+|s|)} dt ds \right), \end{aligned} \quad (2.20)$$

where

$$m_T(k; s, t) = \mathbb{E}_{\mathcal{N}_T} \left[(e^{ik \cdot q_t} - \hat{h}(k))(e^{-ik \cdot q_s} - \hat{h}(k)) \right]. \quad (2.21)$$

We will use two different estimates of this function. One is the uniform bound

$$|m_T(k; s, t)| \leq c_1, \quad (2.22)$$

and the other is

$$|m_T(k; s, t)| \leq |k|^2 \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} p_{s,t}^T(q_1, q_2) (|q_1| + b)(|q_2| + b) d\mathbf{N}^0(q_1) d\mathbf{N}^0(q_2) \quad (2.23)$$

for $|t - s| \geq 1$, where $p_{s,t}^T$ is the probability density with respect to $d\mathbf{N}^0(q_1) d\mathbf{N}^0(q_2)$ of the measure $\mathcal{N}_T$ with $q_t = q_1$, $q_s = q_2$, and $b, c_1 > 0$ are some constants. Using the intrinsic hypercontractivity of the reference measure and the cluster expansion of [4] we have in this case that

$$|p_{s,t}^T(q_1, q_2)| \leq c_2 \quad (2.24)$$

with some $c_2 > 0$, moreover by the exponential fall-off of the stationary measure of the reference process

$$\int_{\mathbb{R}^3} (|q| + b) d\mathbf{N}^0(q) = c_3 < \infty, \quad (2.25)$$

with some $c_3 > 0$. Thus we estimate the exponent in (2.20) by writing

$$\begin{aligned} & \int_{\mathbb{R}^3} dk \int_{-T}^T \int_{-T}^T \frac{|\hat{\rho}(k)|^2}{|k|} m_T(k; s, t) e^{-|k|(|t|+|s|)} ds dt \\ &\leq \int_{\mathbb{R}^3} \frac{|\hat{\rho}(k)|^2}{|k|} \left(c_1 \int_{\substack{-T \leq s, t \leq T \\ |t-s| \leq 1}} e^{-|k|(|t|+|s|)} ds dt + c_2 c_3^2 |k|^2 \int_{\substack{|t|, |s| \leq T \\ |t-s| \geq 1}} e^{-|k|(|t|+|s|)} ds dt \right) dk \\ &\leq \int_{\mathbb{R}^3} \frac{|\hat{\rho}(k)|^2}{|k|} \left(\frac{A_1}{|k|} + A_2 \right) dk < \infty \end{aligned}$$

with some constants $A_1, A_2 > 0$. This then yields a non-zero uniform lower bound on (2.17), and completes the proof of the theorem. $\square$

3 Hamiltonian in the new representation

The Hamiltonian H of the infrared regular representation is defined through (2.15). For having a more concrete expression the natural strategy is to build the Fock space $\mathcal{F}$ over $L^2(\mathcal{S}'(\mathbb{R}^3), dG)$ and to transform H unitarily to an operator acting on $L^2(\mathbb{R}^3, dq) \otimes \mathcal{F}$. We do this in three steps.

First we map $L^2(\mathbb{R}^3, dN^0)$ to $L^2(\mathbb{R}^3, dq)$ through the similarity transformation $R : \phi(q) \mapsto \psi_0(q)\phi(q)$. Thereby the particle Hamiltonian becomes

$$H_p = -\frac{1}{2}\Delta + V(q). \quad (3.1)$$

Next we deal with the Hamiltonian for the free field. First $\mathcal{U}$ defined by (2.5) is used to map $L^2(\mathcal{S}'(\mathbb{R}^3), dG)$ into $L^2(\mathcal{S}'(\mathbb{R}^3), dG_0)$. This map induces a transformation between the shifted, $\mathcal{G}$, and non-shifted, $\mathcal{G}_0$, Gaussian processes, and keeps their generator the same. Then by the Wiener-Itô transform $\mathcal{W} : L^2(\mathcal{S}'(\mathbb{R}^3), dG^0) \rightarrow \mathcal{F}$ this is further mapped into Fock space (for details see e.g. [7]). Finally we transform the interaction. By (2.15) in $L^2(\mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3), dq \times dG)$ the interaction is multiplication by

$$\int_{\mathbb{R}^3} \hat{\varrho}(k) \hat{\xi}(k) \left(e^{ik \cdot q} - \hat{h}(k) \right) e^{-ik \cdot x} dk \quad (3.2)$$

which under $\mathcal{U}$ goes over to the multiplication operator

$$\int_{\mathbb{R}^3} \hat{\varrho}(k) (\hat{\xi}(k) - \hat{\gamma}(k)) \left(e^{ik \cdot q} - \hat{h}(k) \right) e^{-ik \cdot x} dk \quad (3.3)$$

on $L^2(\mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3), dq \times dG_0)$. Consider now

$$\mathcal{V} = (1 \otimes \mathcal{W})(1 \otimes \mathcal{U})(R \otimes 1) \quad (3.4)$$

as an isometry from $L^2(\mathbb{R}^3 \times \mathcal{S}'(\mathbb{R}^3), dP^0)$ to $L^2(\mathbb{R}^3, dq) \otimes \mathcal{F}$. Then

$$\begin{aligned} \mathcal{V} H \mathcal{V}^{-1} &= \left(-\frac{1}{2}\Delta + V(q) \right) \otimes 1 + 1 \otimes \int_{\mathbb{R}^3} |k| a^*(k) a(k) dk \\ &+ \int_{\mathbb{R}^3} \frac{\hat{\varrho}(k)}{\sqrt{2|k|}} \left((e^{ik \cdot q} - \hat{h}(k)) \otimes a(k) + (e^{-ik \cdot q} - \hat{h}(k)) \otimes a^*(k) \right) dk \\ &- \int_{\mathbb{R}^3} \frac{|\hat{\varrho}(k)|^2}{k^2} \hat{h}(k) \left(e^{ik \cdot q} - \hat{h}(k) \right) \otimes 1 dk \\ &\equiv H_N^{\text{ren}}. \end{aligned} \quad (3.5)$$

The standard Nelson Hamiltonian H_N corresponds to setting $\hat{h} \equiv 0$ in (3.5). We see that the subtraction of $\hat{h}$ regularizes the interaction at small k and makes H_N^{ren} a well

defined Hamiltonian having a ground state in $L^2(\mathbb{R}^3, dq) \otimes \mathcal{F}$. In an operator theoretic context, Arai proposed H_N^{ren} of the form (3.5) for the special case $\hat{h} \equiv 1$ and proved that under similar assumptions on the potential as ours H_N^{ren} has a ground state [1].

Obviously, for obtaining the ground state expectation of an operator A of physical interest, one can use H_N^{ren} only in conjunction with the transformation $A \mapsto \mathcal{V}A\mathcal{V}^{-1}$. Thus expectations of A obtained in the standard way by first introducing the cut-off Hamiltonian, taking ground state expectations with it, and subsequently removing the cut-off are identical with computing averages of $\mathcal{V}A\mathcal{V}^{-1}$ by using H_N^{ren} . As proved above, the infrared limit is thus correctly taken care of.

Acknowledgments: R.A.M. thanks Zentrum Mathematik of Technische Universität München for warm hospitality and financial support. He also thanks the Russian Fundamental Research Foundation (grants 99-01-00284 and 00-01-00271) and CRDF (grant NRM 1-2085) for financial support.

References

- [1] Arai, A.: Ground state of the massless Nelson model without infrared cutoff in a non-Fock representation, to appear *Rev. Math. Phys.* (2001)
- [2] Arai, A., Hirokawa, M. and Hiroshima, F.: On the absence of eigenvectors of Hamiltonians in a class of massless quantum field models without infrared cutoff, *J. Funct. Anal.* **168**, 470-497 (1999)
- [3] Fröhlich, J.: On the infrared problem in a model of scalar electrons and massless scalar bosons, *Ann. Inst. Henri Poincaré* **19**, 1-103 (1973)
- [4] Lőrinczi, J. and Minlos, R.A.: Gibbs measures for Brownian paths under the effect of an external and a small pair potential, to appear in *J. Stat. Phys.* (2001)
- [5] Lőrinczi, J., Minlos, R.A. and Spohn, H.: The infrared behaviour in Nelson's model of a quantum particle coupled to a massless scalar field, submitted for publication (2001)
- [6] Nelson, E.: Interaction of nonrelativistic particles with a quantized scalar field, *J. Math. Phys.* **5**, 1190-1197 (1964)
- [7] Obata, N.: *White Noise Calculus and Fock Space*, Springer, 1994